

# Ecological site F002XB002OR

## Levee Group

Last updated: 12/03/2024  
Accessed: 05/20/2025

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 002X–Willamette and Puget Sound Valleys

The Willamette and Puget Sound Valleys Major Land Resource Area (MLRA 2) is located in western Washington and Oregon. It occupies a forearc basin between coast ranges and the Cascade Mountain volcanic arc. The northern part contains Pleistocene drift, outwash, lacustrine and glaciomarine deposits associated with continental glaciers. The southern part contains Late Pleistocene deposits from glacial outburst floods (Missoula Floods). Climate is mild and moist, with a long growing season. Mean annual precipitation ranges from 20 to 60 inches, falling mostly in fall, winter, and spring. Summers are dry. Soil temperature regime is mesic and soil moisture regimes are xeric and aquic.

Most sites in this MLRA can support forested vegetation, but some were maintained as prairie, savanna, or woodland through cultural burning prior to Euro-American settlement. Puget Sound has a moderating effect on temperatures and humidity can be higher in the northern part of the MLRA. Douglas-fir (*Pseudotsuga menziesii*) is widespread throughout. Oregon white oak (*Quercus garryana*) is common on uplands in the south and on warm, exposed or droughty sites in the north. Pacific madrone (*Arbutus menziesii*) occurs in areas close to salt water. Western hemlock (*Tsuga heterophylla*) is codominant with Douglas-fir in the north. Floodplains usually contain black cottonwood (*Populus balsamifera* ssp. *trichocarpa*) and red alder (*Alnus rubra*). Oregon ash (*Fraxinus latifolia*) is typical of forested wetlands in the south. Forestry, urban development, and cultivated agriculture are currently the most extensive land uses (Soil Survey Staff, 2006).

### LRU notes

The Portland Basin and Hills Land Resource Unit (LRU B) is located in northwest Oregon and southwest Washington. It includes the Portland Basin and surrounding hills. Isolated

areas of LRU C (Willamette Valley) occur below 400 feet in the Tualitan Valley on loamy or silty Missoula Flood deposits. The Columbia River Gorge borders this LRU on the east. Brackish tidewater beginning near the town of Cathlamet marks the northwestern limit of this LRU along the Columbia River floodplain. Elevation ranges from sea level to about 1200 feet. Topography is flat to steep. Major landforms include the Columbia River floodplain, glaciofluvial terraces, hills, and foothills. The valley floor is underlain by Pleistocene fluvial deposits (Rowland Formation). Hills and foothills are underlain by Eocene to Pliocene sedimentary rocks (Yamhill, Nestucca, Scotts Mills, Molalla, and Troutdale Formations), Miocene Columbia River Basalt, or Plio-Pleistocene Boring Lavas (Orr et al., 1992). Gravelly or sandy Late Pleistocene Missoula Flood deposits can occur below 400 feet elevation. Hills are covered in loess, and fragipans (brittle subsoil layers) are common.

Mean annual precipitation ranges from 35 to 60 inches. Most falls as rain between October and May. The frost-free period ranges from 160 to 210 days. Ice storms occur each winter. Locations near the Columbia River Gorge experience strong winds. Most locations experience less summer moisture stress compared with the main Willamette Valley; summertime average daily maximum temperatures at Vancouver, WA are 1 to 3 degrees F cooler compared with Corvallis, OR (Agricultural Climate Information System, 2007a, 2007b).

Cultural fire use prior to Euro-American settlement was apparently less than in the main Willamette Valley, though it was used in some areas. General Land Office (GLO) land surveys conducted between 1851 and 1910 indicate that forest and woodland communities were more prevalent than prairies and savannas (Hulse et al., 2002). Forested reference community phases have been chosen for these upland ecological sites.

Presence of Oregon white oak (*Quercus garryana*), and absence of western hemlock (*Tsuga heterophylla*) distinguish this area from coast range (MLRA 1) and Cascade mountain (MLRA 3) ecological types in Oregon. Relative abundance of western redcedar (*Thuja plicata*) helps distinguish this area from the Willamette Valley (LRU C).

## **Classification relationships**

This ecological site group fits within the following LANDFIRE Biophysical Setting (BpS):

- LANDFIRE Biophysical Setting: North Pacific Lowland Riparian Forest and Shrubland (0711560)

## **Ecological site concept**

This site occurs on the Columbia River floodplain. Adjacent river reaches contain freshwater but may be tidally-influenced. Soils are very deep and excessively or somewhat excessively drained. The rooting zone is dry 45 to 70 consecutive days during the summer. Occasional to frequent flooding which scours and deposits sand and gravel,

relatively neutral soil reaction, humid microclimate, and connection to the water table favor the development of deciduous forest.

**Table 1. Dominant plant species**

Tree	(1) <i>Populus balsamifera ssp. trichocarpa</i> (2) <i>Alnus rubra</i>
Shrub	(1) <i>Cornus sericea</i> (2) <i>Physocarpus capitatus</i>
Herbaceous	Not specified

## Physiographic features

Landform: floodplains

Parent material: coarse alluvium

Elevation: 10 to 40 feet

Slope: 0 to 1 percent

Flooding: frequent or occasional; brief duration

Ponding: none

**Table 2. Representative physiographic features**

Landforms	(1) Flood plain
Flooding duration	Brief (2 to 7 days)
Flooding frequency	Occasional to frequent
Ponding frequency	None
Elevation	10–40 ft
Slope	0–1%
Aspect	Aspect is not a significant factor

## Climatic features

Mean annual air temperature: 50 to 54 degrees F

Mean annual precipitation: 35 to 45 inches

Frost free period: 165 to 210 days

## Influencing water features

This site occurs on the Columbia River floodplain. Adjacent river reaches contain freshwater but may be tidally-influenced. Prior to flood control, annual flooding peaked during June and July. The rooting zone is dry 45 to 70 consecutive days during the

summer which is longer than most sites in this LRU. Occasional to frequent flooding which scours and deposits sand and gravel, relatively neutral soil reaction, humid microclimate, and connection to the water table favor the development of deciduous forest.

**Wetland description**

No wetlands associated with this site.

**Soil features**

- Drainage class: excessively or somewhat excessively drained
- Parent material: coarse alluvium
- Soil restrictive feature(s): none
- Soil moisture regime: xeric
- Soil moisture subclass: typic
- Soil temperature regime: mesic
- Particle-size family(s): coarse-loamy or sandy-skeletal
- Soil mineralogy: mixed
- Cation exchange capacity: superactive
- Soil reaction: neutral to moderately acid

Soils in this group are coarse-textured. Gravelly or sandy material exists in the soil profile. The rooting zone for herbaceous plants is dry 45 to 70 consecutive days during the summer which is longer than most sites in this LRU. However, some plants may be able to extract moisture from the water table. Soils classify as Entisols. Organic matter tends to be low.

Soils correlated with this site include Camas, Newberg, and Pilchuck, and Burlington.

**Table 3. Representative soil features**

Parent material	(1) Alluvium
Drainage class	Somewhat excessively drained to excessively drained

**Ecological dynamics**

Central Concept

This site occurs on the Columbia River floodplain. Adjacent river reaches contain freshwater but may be tidally-influenced. Prior to flood control, annual flooding peaked during June and July. Soils are very deep and excessively or somewhat excessively drained. The rooting zone is dry 45 to 70 consecutive days during the summer which is longer than most sites in this LRU. Occasional to frequent flooding which scours and deposits sand and gravel, relatively neutral soil reaction, humid microclimate, and connection to the water table favor the development of deciduous forest. The reference

plant community is black cottonwood - red alder / shrubs / forbs.

## Range in Variability

Dunes correlated with Burlington soil series are included in this site concept but contrasting geomorphology and ecological dynamics may warrant a distinct site.

## Disturbance

High-energy flooding is a primary disturbance process. Prior to flood control, annual floods peaked in June or July (Christy 2004). Since flood control, flooding frequency and sediment supply has decreased. The contemporary flooding regime may be variable across space due to constructed levees and pumping systems. If the associated stream channel moves away from this site, flooding energy diminishes and soils thicken. Natural fire is infrequent and stand-replacing; the estimated mean fire return interval is 750 years (Henderson and Leshner, 2007). Tree-throw occurs in forested communities.

## Plant Composition

Representative native plants are listed below. Not all species are present within the same community phase. Plant lists (especially for grasses, grasslikes, and forbs) are incomplete.

### TREES AND SHRUBS:

willow (*Salix* spp.)

black cottonwood (*Populus balsamifera* ssp. *trichocarpa*)

red alder (*Alnus rubra*)

white alder (*Alnus rhombifolia*)

Oregon ash (*Fraxinus latifolia*)


redosier dogwood (*Cornus sericea*)

Pacific ninebark (*Physocarpus capitatus*)


California blackberry (*Rubus ursinus*)

## State and transition model


### Ecosystem states



### State 1 submodel, plant communities



### State 2 submodel, plant communities



## State 1 Reference

Flooding occurs annually, but high-energy events which rearrange channels, scour, or deposit gravelly alluvium occur less often and are the primary form of disturbance. Invasive plant species are insignificant in this state.

### Community 1.1

#### Deciduous Forest Black Cottonwood, Red Alder, Shrubs, and Forbs

Structure: deciduous forest The overstory consists mainly of black cottonwood or red alder. Ponderosa pine, Douglas-fir, grand fir, or western redcedar may be occasionally present. The shrub layer is deciduous. Willow may be present but is suppressed by shading. Shade-tolerant forbs are present.

### Community 1.2

#### Deciduous Shrub, Field Willows, Grasses, and Forbs

Structure: deciduous shrub field This community is dominated by willow. Black cottonwood or alder saplings are also present, establishing from seed. Black cottonwood can also

establish from twigs buried in moist sediment (Adams, et al. 1987). Grasses and forbs are present.

### **Pathway 1.1A**

#### **Community 1.1 to 1.2**

This pathway represents a high energy flood that destroys the plant community and either exposes or deposits sandy and gravelly material. Soil litter layer is removed.

### **Pathway 1.2B**

#### **Community 1.2 to 1.1**

This pathway represents vegetative growth, entrapment of loamy sediment, and thickening of loamy or silty A horizons. Soil develops a litter layer consisting mainly of leaves.

## **State 2**

### **Converted Invaded**

This state represents post-cultivation conditions with flood control that may best fit within land-use models in future work. Weedy invasive species are usually present and competitive. Hydrology is not altered by draining or filling.

### **Community 2.1**

#### **Cropland or Pastureland**

Structure: annual or perennial crop, tame pasture, or orchard

### **Community 2.2**

#### **Prairie or Shrub Field, Himalayan Blackberry, Pasture Grasses, and Annual Grasses**

Structure: weedy shrub field or prairie This community consists mainly of aggressive weeds, naturalized pasture grasses, or non-native annual grasses. Himalayan blackberry (*Rubus armeniacus*) is aggressive following ground disturbance. Reed canarygrass (*Phalaris arundinacea*) may be present. Forbs such as Canada thistle (*Cirsium arvense*) and bull thistle (*Cirsium vulgare*) may be common. Saplings of native deciduous trees regenerate from seed when imported by flooding.

### **Community 2.3**

#### **Deciduous Forest Black Cottonwood, Alder, Ash, Deciduous Shrubs, Himalayan Blackberry, and Weedy Forbs**

Structure: Closed deciduous forest The overstory consists of an overstory of black

cottonwood, alder, and Oregon ash. The shrub layer is deciduous. The understory has low species diversity and consists of weedy, shade-tolerant shrubs and forbs. Himalayan blackberry may persist under forest canopy.

### **Pathway 2.1A**

#### **Community 2.1 to 2.2**

This pathway represents abandonment. Tillage and other management ceases.

### **Pathway 2.2B**

#### **Community 2.2 to 2.1**

This pathway represents resumed tillage and agricultural management.

### **Pathway 2.2A**

#### **Community 2.2 to 2.3**

This pathway represents continued abandonment and growth over time. Soil develops a litter layer that consists mainly of leaves.

### **Pathway 2.3C**

#### **Community 2.3 to 2.1**

This pathway represents tree and stump removal with resumed tillage and agricultural management.

### **Pathway 2.3B**

#### **Community 2.3 to 2.2**

This pathway represents tree removal alone.

### **Transition T1A**

#### **State 1 to 2**

This pathway represents tillage to the extent that root systems and seed banks of native plants are depleted. Soil litter layer is removed. Invasive plant species are introduced.

### **Restoration pathway R2A**

#### **State 2 to 1**

This pathway represents a high energy flood that destroys the plant community and either exposes or deposits sandy and gravelly material. This pathway requires a migrating stream channel to return to the site.



## **Additional community tables**

### **Other references**

Adams, A. B., Dale, V. H., Smith, E. P., and Kruckeberg, A. R. (1987). Plant survival, growth form and regeneration following the 18 May 1980 eruption of Mount St. Helens, Washington. *Northwest Science*, 61(3): 160-170.  
<http://research.wsulibs.wsu.edu/xmlui/bitstream/handle/2376/1760/v61%20p160%20Adams%20et%20al.PDF?sequence>

Agricultural Climate Information System. (2007). WETS Station Data for Corvallis State University, OR, 1971-2000. [Online]. Available at <http://agacis.rcc-acis.org/?fips=41003> (accessed on 5/7/2020).

Agricultural Climate Information System. (2007). WETS Station Data for Vancouver 4 NNE, WA, 1971-2000. [Online]. Available at <http://agacis.rcc-acis.org/?fips=53011> (accessed on 5/7/2020).

Balster, C.A., and Parsons, R.B. (1968). Geomorphology and soils Willamette Valley, Oregon. Oregon State University Experiment Station Special Report 265.  
<https://ir.library.oregonstate.edu/downloads/mg74qm961>

Christy, J., and Alverson, E. (2011). Historical vegetation of the Willamette Valley, Oregon, circa 1850. *Northwest Science*. 85(2):93-107. <https://doi.org/10.3955/046.085.0202>

Christy, J.A., Alverson, E.R., Dougherty, M.P., Kolar, S.C., Alton, C.W., Hawes, S.M., Ashkenas, L., and Minear, P. (2011). GLO historical vegetation of the Willamette Valley, Oregon, 1851-1910. ArcMap shapefile, Version 2011\_04. Oregon Biodiversity Information Center, Portland State University. Available at [http://www.pdx.edu/sites/www.pdx.edu.pnwlamp/files/glo\\_willamette\\_2011\\_04.zip](http://www.pdx.edu/sites/www.pdx.edu.pnwlamp/files/glo_willamette_2011_04.zip) (accessed on 11/14/2019).

Franklin, J., and Dyrness, C. (1973). Interior valleys of western Oregon. p. 110-129. In *Natural Vegetation of Oregon and Washington*. United States Department of Agriculture Forest Service, Pacific Northwest Forest and Range Experiment Station. General Technical Report PNW-8.

Henderson, J., and Leshner, R. (2007). North Pacific Lowland Riparian Forest and Shrubland. p. 311-314. In *LANDFIRE Biophysical Setting Model Descriptions*. [Online]. Available at [https://www.landfire.gov/national\\_veg\\_models\\_op2.php](https://www.landfire.gov/national_veg_models_op2.php) (accessed on 6/3/2020).

Hulse, D., Gregory, S., and Baker, J. (2002). Presettlement Vegetation circa 1850. p. 38-39. In *Pacific Northwest Ecosystem Research Consortium (ed.) Willamette River Basin Planning Atlas: Trajectories of Environmental and Ecological Change*. [Online]. Available

at

[http://www.fsl.orst.edu/pnwer/wrb/Atlas\\_web\\_compressed/4.Biotic\\_Systems/4b.preserve\\_g\\_web.pdf](http://www.fsl.orst.edu/pnwer/wrb/Atlas_web_compressed/4.Biotic_Systems/4b.preserve_g_web.pdf) (accessed on 9/28/2015).

Johannessen, C. L., Davenport, W.A., Millet, A., and McWilliams, S. (1971). The vegetation of the Willamette Valley. *Annals of the Association of American Geographers*. 61(2):286-302.

Kim, K.D., Ewing, K., and Giblin, D. E. (2006). Controlling *Phalaris arundinacea* (reed canarygrass) with live willow stakes: a density-dependent response. *Ecological Engineering*. 27(3): 219-227.

<http://depts.washington.edu/waipc/docs/Phalaris%20arundinacea.pdf>

Orr, E., Orr, W., and Baldwin, E. (1992). Willamette Valley. p. 203-221. In *Geology of Oregon*. 4th ed. Kendall/Hunt Publishing Company.

Reckendorf, F. (1993). Geomorphology, stratigraphy, and soil interpretations, Willamette Valley, Oregon. p. 178-199. In J.M. Kimble (ed.) *Proceedings of the Eighth International Soil Management Workshop: Utilization of Soil Survey Information for Sustainable Land Use*. Oregon, California, and Nevada. 11-24 July 1992; May 1993. United States Department of Agriculture Soil Conservation Service National Soil Survey Center.

Schoeneberger, P.J., Wysocki, D.A., Benham, E.C., and Soil Survey Staff. (2012). *Field book for describing and sampling soils*, Version 3.0. Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. (2006). *Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin*. Agricultural Handbook 296. [https://www.nrcs.usda.gov/Internet/FSE\\_DOCUMENTS/nrcs142p2\\_050898.pdf](https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_050898.pdf)

Soil Survey Staff. (2014). *Keys to Soil Taxonomy*, 12th ed. USDA-Natural Resources Conservation Service, Washington, DC.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. *Official Soil Series Descriptions*. Online. Available at [https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/geol/?cid=nrcs142p2\\_053587](https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/geol/?cid=nrcs142p2_053587) (accessed 2019 to 2020).

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. *Soil Survey Geographic (SSURGO) Database for Oregon* (multiple counties). [Online]. Available at <https://websoilsurvey.sc.egov.usda.gov/App/HomePage.htm> (accessed in 2020).

Yeats, R.S., Graven, E.P., Werner, K.S., Goldfinger, C. and Popowski, T. (1996).

Tectonics of the Willamette Valley, Oregon. p. 183-222. In Rogers, Albert M., Walsh, Timothy J., Kockelman, William J., and Priest, George R. (ed.) Assessing earthquake hazards and reducing risk in the Pacific Northwest. US Geological Survey Professional Paper 1560.

## Approval

Kirt Walstad, 12/03/2024

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	10/03/2023
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

### 1. Number and extent of rills:

---

### 2. Presence of water flow patterns:

---

### 3. Number and height of erosional pedestals or terracettes:

---

### 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

---

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

17. **Perennial plant reproductive capability:**

---