

Ecological site R007XY988WA

Wetland Complex

Last updated: 2/06/2025
Accessed: 05/21/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 007X–Columbia Basin

This MLRA is in the Walla Walla Plateau section of the Columbia Plateaus province of the Intermontane Plateaus. The Columbia River flows through this MLRA, and the Snake and Yakima Rivers join the Columbia River within it. This MLRA is almost entirely underlain by Miocene basalt flows. Columbia River Basalt is covered in most areas with as much as 200 feet of eolian, lacustrine, and alluvial deposits. The dominant soil orders in this MLRA are Aridisols and Entisols. The soils in the area dominantly have a mesic temperature regime, an aridic moisture regime, and mixed mineralogy. They generally are moderately deep to very deep and well drained to excessively drained.

Classification relationships

Major Land Resource Area (MLRA): 007X – Columbia Basin

LRU – Common Resource Areas (CRA):

- 7.1 – Sandy Missoula Flood Deposits
- 7.2 – Silty Missoula Flood Deposits
- 7.3 – Dry Loess Islands
- 7.4 – Dry Yakima Folds
- 7.5 – Yakima Valley – Pleistocene Lake Basins

Ecological site concept

In the upland setting ecological sites are often expansive, and thus, can be delineated and separated on aerial photos. But in the landscape position of bottoms, basins and

depressions this is rarely the case as small changes in soil chemistry, the water table and elevation or aspect results in significant changes in plant community composition. In short distances there are often big swings of available water holding capacity, and soils can go from hydric to non-hydric, or from saline-sodic to not. So, in bottoms, riparian areas and depressions, ecological sites and community phases occur as small spots, strips and patches, or as narrow rings around vernal ponds. And generally, in a matter of steps one can walk across several ecological sites. On any given site location, two or more of these ecological sites occur as a patchwork – Loamy Bottom, Alkali Terrace, Sodic Flat, Wetland Complex and Riparian Complex. These ecological sites may need to be mapped as a complex when doing resource inventory.

Diagnostics:

Wetland Complex ecological site in MLRAs 007X, 008X, & 009X is a bottomland site, is the “classic wetland” and is characterized by two conditions – hydric soil and aquatic plants. This small patch ecosystem sits on the lowest position of the landscape, on landforms such as depressions, bottoms, floodplains and basins. Wetland Complex also occurs on pond and lake fringes, and along slow-moving streams and rivers. These sites are so small they are indicated on a soil map as a spot symbol.

Wetland Complex ecological site is part of the lentic (standing water) ecosystem. Wetlands are frequently or continually inundated by up to two feet of water. Water level fluctuations support the development of different wetland zones (floating, submergent, emergent). This ecological site only considers the emergent vegetative zone of the wetland (where plants rise above the water surface). The floating and submergent zones are not considered in this description.

Soils are saturated to the surface or there is standing water for an extended portion of the growing season. Thus, the soils show all the signs of hydric soils such as mottling and greying. These saturated wetland soils are not saline or sodic but, are hydric. The soils are moderately deep to deep, silt loam or sandy loam texture.

These plant communities are exclusively herbaceous (non-woody) and predominately wetland obligate species. Cattails, bulrush, sedges, wetland grasses and Baltic rush are major species. Wetlands often have low species diversity as many of the dominant species form dense monocultures. Wetland Complex remains wet all season and rarely, if ever, burn.

A subset of this ecological site occurs around the edge of basalt pothole ponds. In addition to the herbaceous species, this subset can have woody species such as aspen, coyote willow, wood rose and hawthorn.

Principle Vegetative Drivers:

Prolonged saturated and anaerobic soil conditions drive the vegetative expression of the Wetland Complex ecological site. Seasonal fluctuations in water levels control vegetation patterns. This site is dominated by hydrophytic species.

Associated sites

R007XY930WA	Loamy Bottom
R007XY970WA	Alkali Terrace
R007XY978WA	Sodic Flat
R007XY720WA	Riparian Complex
R007XY130WA	Loamy
R007XY120WA	Stony
R007XY153WA	Cool Loamy

Similar sites

R008XY988WA	Wetland Complex
R009XY988WA	Wetland Complex

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Beckmannia syzigachne</i> (2) <i>Schoenoplectus acutus</i>

Physiographic features

The landscape is part of the Columbia basalt plateau. This ecological site sits on the lowest position on the landscape on landforms such as bottoms, floodplains, basins & depressions. Wetland complex also occurs as fringes around ponds and lakes at elevations of 300 to 1,500 feet. In bottoms, riparian areas and depressions, ecological sites and community phases occur as small spots, strips and patches, or as narrow rings around vernal ponds.

Physiographic Division: Intermontane Plateau

Physiographic Province: Columbia Plateau

Physiographic Sections: Walla Walla Plateau Section

Table 2. Representative physiographic features

Landforms	(1) Hills (2) Plateau (3) Flood plain (4) Depression (5) Basin floor
Flooding frequency	None to occasional
Ponding frequency	None
Elevation	91–457 m
Slope	0–3%
Water table depth	0–30 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	Not specified
Slope	0–5%
Water table depth	Not specified

Climatic features

The climate is characterized by moderately cold, wet winters, and hot, dry summers, with limited precipitation due to the rain shadow effect of the Cascades. This MLRA is the warmest and driest MLRA within the Columbia Plateau geographic area. Seventy to seventy-five percent of the precipitation comes late-October through March as a mixture of rain and snow. Precipitation that comes after March is not as effective for plant growth, but June through early-October can be dry. Freezing temperatures generally occur from late-October through early-April. Temperature extremes are -10 degrees Fahrenheit in winter and 110 degrees Fahrenheit in summer.

Table 4. Representative climatic features

Frost-free period (characteristic range)	150-180 days
Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	152-254 mm
Frost-free period (actual range)	140-200 days
Freeze-free period (actual range)	
Precipitation total (actual range)	

Influencing water features

A plant's ability to grow on a site and overall plant production is determined by soil-water-plant relationships:

1. Whether rain and melting snow run off-site or infiltrate into the soil
2. Whether soil condition remain aerobic or become saturated and anaerobic
3. How quickly the soil reaches the wilting point

Water is at or above the surface for most of the growing season. Seasonal flooding, runoff and discharging groundwater maintain saturated and anaerobic soil conditions.

Soil features

Dominantly wetlands occur as spot symbols on soil map there isn't much soils information for them. What we do have for this ecological site soil components are dominantly Fluvaquents and Psammaquents great groups of the Entisol taxonomic order. Soils in the Histosols taxonomic order should exist in this ecological site, but a map unit hasn't been large enough to map. Soils are deep and very deep. Average available water capacity of about 18 inches (45.7 cm) in the 0 to 40 inches (0 to 100 cm) depth range.

Soil parent material is dominantly alluvium derived from mixed sources with possibly minor amounts of ash in the upper part of the soil.

The associated soils are Wanser, Fluvaquents and similar soils.

Table 5. Representative soil features

Parent material	(1) Alluvium
Surface texture	(1) Silt loam (2) Fine sand
Family particle size	(1) Coarse-silty (2) Sandy or sandy-skeletal
Drainage class	Very poorly drained to poorly drained
Depth to restrictive layer	51–152 cm
Soil depth	152 cm
Surface fragment cover <=3"	1%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	45.72 cm
Calcium carbonate equivalent (Depth not specified)	0%

Electrical conductivity (Depth not specified)	0 mmhos/cm
Sodium adsorption ratio (Depth not specified)	0
Soil reaction (1:1 water) (0-25.4cm)	6.6–7.9
Subsurface fragment volume <=3" (Depth not specified)	5%
Subsurface fragment volume >3" (Depth not specified)	2%

Table 6. Representative soil features (actual values)

Drainage class	Not specified
Depth to restrictive layer	Not specified
Soil depth	Not specified
Surface fragment cover <=3"	0–5%
Surface fragment cover >3"	0–5%
Available water capacity (0-101.6cm)	30.48–60.96 cm
Calcium carbonate equivalent (Depth not specified)	Not specified
Electrical conductivity (Depth not specified)	Not specified
Sodium adsorption ratio (Depth not specified)	Not specified
Soil reaction (1:1 water) (0-25.4cm)	Not specified
Subsurface fragment volume <=3" (Depth not specified)	0–15%
Subsurface fragment volume >3" (Depth not specified)	0–10%

Ecological dynamics

The Wetland Complex ecological site in MLRA 007X produces about 10,000 pounds per acre of biomass annually.

Almost all wetlands have been impacted by widespread degradation from (1) hydrologic alteration, (2) invasion by invasive weeds such as reed canarygrass, or (3) excessive

grazing. Many wetlands are completely dominated by invasive species.

Cattails are aquatic, perennial plants found in a variety of wetland habitats. These are often the first wetlands plants to colonize areas of newly exposed wet mud, with their abundant wind-dispersed seeds. Cattails also spread by rhizomes, forming large, interconnected stands.

Hardstem bulrush is a perennial, heavily rhizomatous wetland plant. It forms large stands with young plants on the outside and the older plants toward the center. It is generally found in areas of standing water ranging from 4 inches to 6 feet in depth but does not tolerate long periods of deep water.

Reed canarygrass is a circumboreal species, native to north-temperate regions. It grows in wet areas such as edges of lakes, ponds, ditches and creeks, often forming dense stands, in some areas it is a problematic weed. North American populations may be a mix of native strains, European strains and agronomic cultivars. Reed canarygrass frequents saturated soils but, cannot survive extended periods of standing water. Rated FACW 67 to 99 percent occurrence in wetlands.

American mannagrass is a perennial wetland plant that approaches six feet in height. American sloughgrass is an annual or short-lived perennial.

Nebraska sedge grows in wetlands across central and western US. It tolerates submersion for long periods and, also, alkaline conditions. Nebraska sedge has bluish leaves and produces a dense network of rhizomes.

Most freshwater marshes and wetlands experience seasonal and episodic flooding. Water level fluctuations support the development of different marsh zones. Seasonal fluctuations in water levels control vegetation patterns and invertebrate communities. Often the herbaceous community of the Wetland Complex ecological site has low species diversity as many of the dominant species form dense monocultures.

Wetlands almost never burn and because of standing water receives limited grazing pressure.

In Washington, wetland communities in a sagebrush steppe ecosystem provide habitat for a variety of wetland and upland wildlife species.

State and transition model

Reference State

Community Phase 1.1

Complex or monoculture of Native Species
Native Sedges, Rushes, Forbs, Grasses

T1

State 2 – Altered State

Community Phase 2.1

Intensive Disturbance
Wetland drained or filled
Vegetation displaced by invasive species
Vegetation killed
Excessive grazing

State 1 Reference

State 1 represents stable wetlands with no invasive or exotic weed species. Often this site has low species diversity as dominant species can form dense monocultures. Reference Community 1.1 can be dominated by cattails or, bulrush or, Nebraska sedge or, American mannagrass or, Baltic rush. At-risk Communities: All communities in the reference state

are at risk because of heavy grazing pressure and other human manipulations to meadows. Reference Community 1.1 for Wetland Complex ecological site Percentages for plant species composition below are an approximated weight. The composition of pristine sites can vary somewhat due to variations in site conditions. Wet Meadow species that are OBLIGATE went to this site Often this site has low species diversity as many of the dominant species form dense monocultures. Some areas are 100 percent cattails, others 100 percent bulrush, and other areas 100 percent Nebraska sedge, etc. Native Grasses: BESY American sloughgrass GLCR American mannagrass GLST fowl mannagrass Native Sedges: CAUT Northwest Territory sedge CANE2 Nebraska sedge CAPE42 woolly sedge Native rushes: SCAC hardstem bulrush SCTA softstem bulrush SCMI paniced bulrush JUBA Baltic rush Native forbs: TYLA cattails POLYG smartweed Pathways within State 1 (Reference State) None as there is only one community in Reference State.

Community 1.1

Complex or Monoculture of Native Sedges, Rushes, Forbs, and Grasses

This site often has low diversity as many of the dominant species form dense monocultures.

Dominant plant species

- American sloughgrass (*Beckmannia syzigachne*), grass
- American mannagrass (*Glyceria grandis*), grass
- fowl mannagrass (*Glyceria striata*), grass
- Northwest Territory sedge (*Carex utriculata*), other herbaceous
- Nebraska sedge (*Carex nebrascensis*), other herbaceous
- woolly sedge (*Carex pellita*), other herbaceous
- hardstem bulrush (*Schoenoplectus acutus*), other herbaceous
- softstem bulrush (*Schoenoplectus tabernaemontani*), other herbaceous
- paniced bulrush (*Scirpus microcarpus*), other herbaceous
- arctic rush (*Juncus arcticus*), other herbaceous
- broadleaf cattail (*Typha latifolia*), other herbaceous
- knotweed (*Polygonum*), other herbaceous

State 2

Altered

State 2 represents an altered state because of intensive disturbance. The wetland may have been drained or filled to convert to a different land use. Or, the wetland may have experienced excessive grazing. Invasive species such as phragmites, reed canarygrass, and purple loosestrife may dominate the plant community in the altered state. Community Phases for State 2: Can have several variations: Reed canarygrass Purple loosestrife Phragmites

Community 2.1

Invasive Species

Intensive disturbance. Wetland may be drained or filled. Vegetation is displaced by invasive species.

Transition T1A

State 1 to 2

Result: Transition from Reference State to altered State 2 Ecological process: Wetland hydrology altered, and site may no longer have wetland functions. Invasive species colonize the site and over time dominate the stand. Primary Trigger: both deliberate and unintentional, human-caused, alterations such as drainage, filling the wetland with soil, herbicide drift, deliberate use of herbicides or grazing pressure. Indicators: occurrence of invasive species where there has been none. Declining cover of native species and increasing cover of invasive species. Site is much drier than previously.

Additional community tables

Inventory data references

Data to populate Reference Community came from several sources: (1) NRCS ecological sites from 2004, (2) Soil Conservation Service range sites from 1980s and 1990s, (3) Daubenmire's habitat types, and (4) ecological systems from Natural Heritage Program

Other references

Boling M., Frazier B., Busacca, A., General Soil Map of Washington, Washington State University, 1998

Crawford, Rex C., Riparian Vegetation Classification of the Columbia Basin, Washington, March 2003.

Environmental Protection Agency, map of Level III and IV Ecoregions of Washington, June 2010

Natural Resources Conservation Service, map of Common Resource Areas of Washington, 2003

Rocchio, Joseph & Crawford, Rex C., Ecological Systems of Washington State. A Guide to Identification. Washington State Department of Natural Resources, October 2015. Pages 156-161 Inter-Mountain Basin Big Sagebrush.

Rouse, Gerald, MLRA 8 Ecological Sites as referenced from Natural Resources Conservation Service-Washington FOTG, 2004

Soil Conservation Service, Range Sites for MLRA 8 from 1980s and 1990s

Contributors

K. Guinn
R. Fleenor
W. Keller
K. Bomberger
K. Paup-Lefferts

Approval

Kirt Walstad, 2/06/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	02/01/2025
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-