

Ecological site R009XY015OR Clayey 14-17 PZ

Last updated: 5/05/2025 Accessed: 05/22/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Jeff Repp
Contact for lead author	Oregon NRCS State Rangeland Management Specialist
Date	07/30/2012
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1.	Number and extent of rills: None, slight to moderate sheet & rill erosion hazard
2.	Presence of water flow patterns: None

3. Number and height of erosional pedestals or terracettes: None to some (<1.0")

moss, plant canopy are not bare ground): 5-15%
Number of gullies and erosion associated with gullies: None
Extent of wind scoured, blowouts and/or depositional areas: None, slight wind erosion hazard
Amount of litter movement (describe size and distance expected to travel): Fine - limited movement
Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Significantly resistant to erosion; aggregate stability = 3
Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Moderately deep (with areas of rock outcrop), moderately well drained to well drained with silty clay loam to silt loam surfaces with up to 35% cobbles or stones; moderate OM (2-3%)
Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Moderate ground cover (60-70%) and gentle slopes (0-12% may be up to 20%) moderately limits rainfall impact and overland flow
Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None

	than, greater than, and equal to):
	Dominant: Idaho fescue > Bluebunch wheatgrass > other perennial grasses = dominant forbs > other forbs > Western Juniper
	Sub-dominant:
	Other:
	Additional:
13.	Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Normal decadence and mortality expected
14.	Average percent litter cover (%) and depth (in):
15.	Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): Favorable: 1200, Normal: 700, Unfavorable: 400 lbs/acre/year at high RSI (HCPC)
16.	Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Annual bromes and Medusahead invade sites that have lost deep rooted perennial grass functional groups. Excessive erosion may occur, deteriorating site potential.
17.	Perennial plant reproductive capability: All species should be capable of reproducing annually