

Ecological site R010XA023OR Juniper Shrubby Lava Blisters 10-12 PZ

Last updated: 3/26/2025 Accessed: 05/20/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Jeff Repp and Bruce Franssen
Contact for lead author	State Rangeland Management Specialist for NRCS - Oregon
Date	04/24/2003
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills: None to some.
- 2. Presence of water flow patterns: None to some.
- 3. Number and height of erosional pedestals or terracettes: None.

- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): 10 20%
- 5. Number of gullies and erosion associated with gullies: None.
- 6. Extent of wind scoured, blowouts and/or depositional areas: None to some.
- 7. Amount of litter movement (describe size and distance expected to travel): Fine limited movement.
- Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values): Moderately to slightly resistant to erosion: aggregate stability = 2-4.
- Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Shallow and very shallow, well drained, sandy loams and stony loamy sands dominated by volcanic ash: Weak thin and medium platy to single grain structure, dry color value of 5, 3-4 inches thick; Low OM (1-2%).
- Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Moderate ground cover (40-50%) and slight to severe slopes (0-15% with some as high as 60% on sides of blisters) moderately limit rainfall impact and overland flow.
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None.
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Perennail, deep-rooted bunch-grasses

Sub-dominant: Evergreen shrubs = deciduous shrubs

Other: Evergreen trees = perennial forbs > other shrubs

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Normal decadence and mortality expected.
- 14. Average percent litter cover (%) and depth (in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): Favorable: 900, Normal: 700, Unfavorable: 500 lbs/acre/year.
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Cheatgrass and Medusahead invade sites that have lost deep rooted perennial grass functional groups.
- 17. **Perennial plant reproductive capability:** All species should be capable of reproducing annually.