

Ecological site R013XY037ID Loamy 13-16 PZ ARTRW8/POA

Last updated: 2/13/2025
Accessed: 05/20/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Dave Franzen and Jacy Gibbs Intermountain Range Consultants 17700 Fargo Rd. Wilder, ID 83676
Contact for lead author	Brendan Brazee, State Rangeland Management Specialist USDA- NRCS 9173 W. Barnes Drive, Suite C, Boise, ID 83709
Date	06/24/2009
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** rills can occur on this site. If rills are present they are likely to occur on slopes over 15 percent and immediately following wildfire. They are most likely to occur on silt loam surface textures.
-

2. **Presence of water flow patterns:** water-flow patterns can occur on this site, particularly on

slopes greater than 15 percent. If they occur, they are short and disrupted flows. They are disrupted by cool season grasses and tall shrubs and are not extensive.

3. **Number and height of erosional pedestals or terracettes:** both can occur on this site. Where flow patterns and/or rills are present, a few pedestals may be expected.
-

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** data is not available. On sites in mid-seral status bare ground may range from 50-65 percent.
-

5. **Number of gullies and erosion associated with gullies:** none.
-

6. **Extent of wind scoured, blowouts and/or depositional areas:** usually not present. Immediately following wildfire some soil movement may occur on lighter textured soils. Where sagebrush has repopulated the site after a fire, remnants of past wind scour may be present.
-

7. **Amount of litter movement (describe size and distance expected to travel):** fine litter in the interspaces may move up to 2 feet following a significant run-off event. Coarse litter generally does not move.
-

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** values should range from 3 to 5 but needs to be tested.
-

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** structure ranges from weak very fine granular to weak fine granular structure. Soil organic matter (SOM) generally ranges from 1 to 2 percent. The surface color ranges from dark brown to dark yellowish brown.
-

10. **Effect of community phase composition (relative proportion of different functional**

groups) and spatial distribution on infiltration and runoff: bunchgrasses, especially deep-rooted perennials, slow run-off and increase infiltration. Tall shrubs can catch snow in the interspaces.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** is not present.
-

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: cool season deep -rooted perennial bunchgrasses

Sub-dominant: tall shrubs

Other: perennial forbs

Additional: shallow rooted grasses

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Wyoming big sagebrush will become decadent in the absence of normal fire frequency. Grass and forb mortality will occur as tall shrubs increase.
-

14. **Average percent litter cover (%) and depth (in):** additional litter cover data is needed but is expected to be 5-20 percent to a depth of 0.1 inches. Under mature shrubs litter is <0.5 inches deep and is 90-100 percent ground cover.
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** is 700 pounds per acre (784 kilograms per hectare) in a year with normal temperatures and precipitation. Perennial grasses produce 40-50 percent of the total production, forbs 20-30 percent, and shrubs 25-35 percent.
-

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: includes cheatgrass, *Vulpia* sp., annual mustards, bulbous bluegrass, and spotted and diffuse knapweeds.
-

17. **Perennial plant reproductive capability:** all functional groups have the potential to reproduce in most years.
-