

Ecological site R027XY068NV GRANITIC CLAYPAN 8-10 P.Z.

Last updated: 6/03/2024
Accessed: 05/21/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Patti Novak-Echenique
Contact for lead author	State Rangeland Management Specialist
Date	07/19/2013
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:** Rills are none to rare. A few can be expected on steeper slopes in areas subjected to summer convection storms or rapid spring snowmelt.

- 2. Presence of water flow patterns:** Water flow patterns are none to rare but can be expected in areas subjected to summer convection storms or rapid snowmelt.

- 3. Number and height of erosional pedestals or terracettes:** Pedestals are rare with occurrence typically limited to area within water flow patterns. Frost heaving of shallow rooted

plants should not be considered as normal condition.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare Ground up to 30% depending on amount of surface rock fragments
-

5. **Number of gullies and erosion associated with gullies:** None
-

6. **Extent of wind scoured, blowouts and/or depositional areas:** None
-

7. **Amount of litter movement (describe size and distance expected to travel):** Fine litter (foliage from grasses and annual & perennial forbs) expected to move distance of slope length during intense summer convection storms or rapid snowmelt events. Persistent litter (large woody material) will remain in place except during large rainfall events.
-

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values should be 3 to 6 on most soil textures found on this site. (To be field tested.)
-

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Surface structure is typically prismatic to columnar or subangular blocky. Soil surface colors are brownish grays and soils are typified by an ochric epipedon. Organic matter of the surface 2 to 3 inches is typically 1 to 1.5 percent dropping off quickly below. Organic matter content can be more or less depending on micro-topography.
-

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Perennial herbaceous plants slow runoff and increase infiltration. Shrub canopy and associated litter break raindrop impact and provide opportunity for snow catch and accumulation on site.
-

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** Compacted layers are none. Subsoil argillic horizons are not to be interpreted as compacted.
-

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: deep-rooted, cool season perennial bunchgrasses

Sub-dominant: evergreen shrub (Lahontan sagebrush) >associated shrubs>deep-rooted, cool season, perennial forbs> fibrous, shallow-rooted, cool season, annual and perennial forbs= deep-rooted, cool season, perennial grasses

Other:

Additional: After wildfires, deep-rooted, cool season perennial bunchgrasses, perennial forbs and sprouting shrubs (spiny hopsage, ephedra, rabbitbrush) dominate. Lahontan sagebrush is removed for 5 to 10 years.

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Dead branches within individual shrubs common and standing dead shrub canopy material may be as much as 25% of total woody canopy; some of the mature bunchgrasses (<20%) have dead centers.
-

14. **Average percent litter cover (%) and depth (in):** Between plant interspaces 20-30% and depth <1/4-in.
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** For normal or average growing season, ±600 lbs/ac; Favorable years ±900 lbs/ac and unfavorable years ±350 lbs/ac.
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment**

and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is **NOT** expected in the reference state for the ecological site: Potential invaders include cheatgrass, medusa head, halogeton, Russian thistle and annual mustards.

17. **Perennial plant reproductive capability:** All functional groups should reproduce in average (or normal) and above average growing season year. Reduced growth and reproduction occur during extended or extreme drought conditions.
-