

Ecological site EX043B23A142

Saline Subirrigated (SS) Absaroka Lower Foothills

Last updated: 4/30/2024

Accessed: 05/22/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Ray Gullion, E. Bainter
Contact for lead author	ray.gullion@wy.usda.gov or 307-347-2456
Date	05/01/2008
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Rare to nonexistent.
-

2. **Presence of water flow patterns:** Barely observable.
-

3. **Number and height of erosional pedestals or terracettes:** Rare to nonexistent.
-

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen,**

moss, plant canopy are not bare ground): Bare ground can range from 0-5%.

5. **Number of gullies and erosion associated with gullies:** Active gullies should not be present.

6. **Extent of wind scoured, blowouts and/or depositional areas:** Minimal to nonexistent.

7. **Amount of litter movement (describe size and distance expected to travel):** Herbaceous litter not expected to move.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil Stability Index ratings range from 3 (interspaces) to 6 (under plant canopy), but average values should be 4.0 or greater.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Typically an A-horizon of greater than 2 inches (5 cm) with massive structure and color hues of 10YR or 2.5Y, values of 5-6, and chromas of 2-3. Organic matter typically is 1-2%.

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Plant community consists of 75-90% grasses, 10% forbs, and 0-15% shrubs. Dense plant canopy (80-100%) and litter plus moderate infiltration rates result in minimal to nonexistent runoff. Basal cover is typically greater than 5% for this site and effectively reduces runoff on this site.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction layer exists.

12. **Functional/Structural Groups (list in order of descending dominance by above-ground**

annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: warm season bunchgrasses

Sub-dominant: tall, cool season bunchgrasses

Other: perennial forbs = mid-size, cool season bunchgrasses = cool season rhizomatous grasses = warm season rhizomatous grasses perennial shrubs

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Minimal decadence.
-

14. **Average percent litter cover (%) and depth (in):** Litter ranges from 0-20% of total canopy measurement with total litter (including beneath the plant canopy) from 80-100% expected. Herbaceous litter depth typically ranges from 10-25 mm.
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** English: 2800-3600 lb/ac (3200 lb/ac average); Metric: 3136-4032 kg/ha (3584 kg/ha average).
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Bare ground greater than 20% is the most common indicator of a threshold being crossed. Greasewood and inland saltgrass are common increasers. Russian olive, saltcedar, and foxtail barley may invade the site.
-

17. **Perennial plant reproductive capability:** All species are capable of reproducing, except in

drought years.
