

## **Ecological site R047XA456UT Mountain Stony Loam (antelope bitterbrush)**

Last updated: 2/05/2025  
Accessed: 05/21/2025

---

### **Rangeland health reference sheet**

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	V. Keith Wadman (NRCS Retired).
Contact for lead author	shane.green@ut.usda.gov
Date	11/13/2012
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### **Indicators**

1. **Number and extent of rills:** Rare to Slight. Slight rill development may occur on steeper slopes (> 20%) or on areas located below exposed bedrock or other water shedding areas where increased runoff may occur. Where these rills are present, they should be fairly short (4-8 feet), < 1 inch deep and somewhat widely spaced (5-10 feet). Minor rill development may be observed on all slopes following major thunderstorm or spring runoff events but should heal during the next growing season.
- 
2. **Presence of water flow patterns:** Slight. Some minor evidence of water flow patterns may be found winding around perennial plant bases. They show little evidence of current erosion. They are expected to be short (3-6 feet), stable, sinuous and normally not connected. There

may be very minor evidence of deposition. Evidence of water flow may increase somewhat on slopes > 20%.

---

3. **Number and height of erosional pedestals or terracettes:** Perennial vegetation shows little evidence of erosional pedestalling (1 to 2% of individual plants). Plant roots are covered and most litter remains in place around plant crowns. Terracettes should be absent or, if present, stable. A slight increase in both pedestal and terracette development may occur with increasing slope.
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground ranges from 20% - 25%. Soil surface may be covered by 20 to 70% coarse fragments. Bare ground openings should not be greater than 1 foot in diameter and should normally not be connected.
5. **Number of gullies and erosion associated with gullies:** None to Rare at site level. Scattered landscape level gully channels, however, are a normal component of basin/range environments. Where landscape gullies are present, they should be stable, partially vegetated on their sides and bottoms, with no evidence of head-cutting. Some slight increase in disturbance may be evident following significant weather events or when gullies convey considerable runoff from higher elevation rocky or naturally eroding areas.
6. **Extent of wind scoured, blowouts and/or depositional areas:** None. No evidence of wind generated soil movement is present. Wind caused blowouts and deposition are not present.
7. **Amount of litter movement (describe size and distance expected to travel):** Most litter resides in place with some redistribution caused by water movement. Minor litter removal may occur in flow channels with deposition occurring within 1 to 2 feet at points of obstruction. The majority of litter accumulates at the base of plants. Some grass leaves and small twigs (grass stems) may accumulate in soil depressions adjacent to plants. Woody stems are not likely to move. However, some litter movement is expected (up to 6 feet) with increases in slopes >20% and/or increased runoff resulting from heavy thunderstorms.
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most**

**sites will show a range of values):** This site should have a soil stability rating of 5 or 6 under the plant canopies, and a rating of 4 to 5 in the interspaces. The average rating should be a 5. Soil surface textures are typically loams, very fine sandy loams and silt loams.

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** (Horrocks) Soil surface 0-5 inches. Texture is a very cobbly sandy clay loam; color is very dark grayish brown (10YR 3/2); and structure is weak medium granular. Mollic epipedon ranges to 14 inches. Use the specific information for the soil you are assessing found in the published soil survey to supplement this description.
- 

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Perennial vegetation produces sufficient cover and spatial arrangement to intercept most raindrops and reduce raindrop splash erosion. Litter on soil surface and cryptogamic crusting, where present, also protect the soil surface from splash erosion and encourage higher infiltration. Bare spaces are expected to be small and irregular in shape and usually not connected. Vegetative structure and distribution are usually adequate to capture snow and ensure that snowmelt occurs in a controlled manner, allowing maximum time for infiltration, and reducing runoff and erosion in all but the most extreme storm events. When perennial grasses and shrubs decrease due to natural events such as long-term drought, insect damage, etc., runoff is likely to increase and infiltration be reduced.
- 

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None. Some soils may have natural textural variability within their profiles, including changes in clay content, these should not be mistaken for a compaction pan.
- 

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Sprouting shrubs (bitterbrush, alderleaf mountain mahogany) > Non-sprouting shrub (mountain big sagebrush) = > Perennial bunchgrasses (bluebunch wheatgrass, Letterman needlegrass) > Rhizomatous Grasses (slender wheatgrass).

Sub-dominant: Perennial forbs (arrowleaf balsamroot)

Other: A wide variety of other perennial grasses and both perennial and annual forbs can be expected to occur in the plant community.

Additional: Natural disturbance regimes include fire, drought, and insects. Assumed fire cycle of 40 to 60+ years. Functional/structural groups may appropriately contain non-native species if their ecological function is the same as the native species in the reference. Following a disturbance such as fire, drought, rodents or insects that remove woody vegetation, forbs and perennial grasses (herbaceous species) may dominate the community for a period of time. If a disturbance has not occurred for an extended period of time, woody species may continue to increase. These conditions would reflect different functional community phases within the reference state.

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Slight decadence in the principle shrubs could occur near the end of the fire cycle or during periods of extended drought, or insect infestations. In general, a mix of age classes should be expected with some dead and decadent plants present.
- 

14. **Average percent litter cover (%) and depth ( in):** Litter cover will be heavier under plants. Most litter will be herbaceous and depths of 1 to 2 inches would be considered normal. Perennial vegetation should be well distributed on the site.
- 

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Annual production in air-dry herbage should be approximately 1400 - 1500 #/acre on an average year but could range from 750 - 1750 #/acre during periods of prolonged drought or above average precipitation.
- 

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing**

**what is NOT expected in the reference state for the ecological site:** Cheatgrass, alyssum, mustard species, Canada thistle, black medic, Utah juniper, Gamble oak.

---

17. **Perennial plant reproductive capability:** All perennial plants should have the ability to reproduce in all years, except in extreme drought years. Green rabbitbrush sprouts vigorously following fire. There are no restrictions on either seed or vegetative reproduction. Some seedling recruitment of major species is expected to be present during average and above average growing years.
-