

Ecological site R055BY066ND Thin Claypan

Last updated: 4/25/2025

Accessed: 05/21/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	USDA-NRCS North Dakota
Contact for lead author	NRCS State Rangeland Management Specialist
Date	04/25/2025
Approved by	Suzanne Mayne-Kinney
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Rills are not expected on this site.
-

2. **Presence of water flow patterns:** Water flow patterns are not visible.
-

3. **Number and height of erosional pedestals or terracettes:** Neither pedestals nor terracettes are expected on this site.
-

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground ranges from 10 to 35%. Bare ground patches (including slickspots) should be small (4 to 8 inches in diameter) and disconnected. Animal activity (burrows and ant mounds) may occasionally result in isolated bare patches of up to 24 inches in diameter.
-
5. **Number of gullies and erosion associated with gullies:** Active gullies are not expected on this site.
-
6. **Extent of wind scoured, blowouts and/or depositional areas:** No wind-scoured or depositional areas expected on this site.
-
7. **Amount of litter movement (describe size and distance expected to travel):** Fine/small class of plant litter associated with slick spots may be moved 4 to 8 inches following rain events. Small accumulations of plant litter may be visible.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Stability class averages 3 or greater. Lower averages expected on the "slickspots" or plant interspaces while higher averages expected under plant canopy.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Use soil series description for depth, color, and structure of A-horizon.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Mid- and short-statured rhizomatous grasses and mid- and short-statured bunchgrasses are dominant and well distributed across the site. Forbs are subdominant.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction layer

would be expected except for the naturally occurring pan within 6 inches of the soil surface.

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Phase 1.1:

Mid & short C3 rhizomatous grasses (1); Mid & short C4 bunch grasses (2)

Sub-dominant: Phase 1.1:

Mid & short C3 bunch grasses (2); Forbs (6)

Other: Phase 1.1:

Mid & short C4 rhizomatous grasses; Grass-likes; Shrub

Additional: Due to differences in phenology, root morphology, soil biology relationships, and nutrient cycling Kentucky bluegrass, smooth brome, and crested wheatgrass are included in a new Functional/structural group, mid- and short-statured early cool-season grasses (MSeC3), not expected for this site.

To see a full version 5 rangeland health worksheet with functional/structural group tables, please use the following hyperlink:

https://efotg.sc.egov.usda.gov/references/public/ND/55B_Thin_Claypan_Narrative_FIN

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Rare to not occurring on this site.
-

14. **Average percent litter cover (%) and depth (in):** Plant litter cover is 35 to 55% with a depth of 0.1 to 0.25 inches. Litter is in contact with soil surface.
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Annual air-dry production is 1300 lbs./ac (reference value) with normal precipitation and temperatures. Low and high production years should yield 800 lbs./ac to 1800 lbs./ac, respectively.
-

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: State and local noxious species, Kentucky bluegrass, smooth brome grass, crested wheatgrass, and Eastern red cedar/juniper.
-

17. **Perennial plant reproductive capability:** Noninvasive species in all functional/structural groups are vigorous and capable of reproducing annually under normal weather conditions.
-