

## Ecological site R102AY016SD Very Shallow

Last updated: 5/05/2025 Accessed: 05/20/2025

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	David Schmidt, Tim Nordquist, Stan Boltz
Contact for lead author	
Date	12/04/2007
Approved by	Suzanne Mayne-Kinney
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## **Indicators**

1.	Number and extent of rills: Rills should not be present.
2.	Presence of water flow patterns: Typically not observable.
3.	Number and height of erosional pedestals or terracettes: None.

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen,

	moss, plant canopy are not bare groundy. Bare ground 20-40 %.
5.	Number of gullies and erosion associated with gullies: Active gullies should not be present.
6.	Extent of wind scoured, blowouts and/or depositional areas: None.
7.	Amount of litter movement (describe size and distance expected to travel): Little to no plant litter movement.
8.	Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Stability class usually 4-6. Moderately high root content. Soil surface is resistant to erosion, in large part due to high rock/gravel content.
9.	Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Use soil series description for depth and color of A-horizon.
10.	Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Healthy, native grasses enhance infiltration and reduce runoff.
11.	Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): No compaction layer should be evident.
12.	Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Mid cool-season bunch grass > short warm-season grass

	Sub-dominant: > short cool-season grass > mid warm-season rhizomatous grass > mid warm-season bunch grass = forb > shrub
	Other:
	Additional:
13.	Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Very little to no evidence of decadence or mortality.
14.	Average percent litter cover (%) and depth (in): 20-40%, less than 0.5 inch thick. Litter cover is in contact with soil surface.
15.	Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): 1600 – 2200 lbs./acre air-dry weight, average 1,900 lbs./acre air-dry weight
16.	Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Refer to State and Local Noxious Weed List
17.	Perennial plant reproductive capability: All species are capable of reproducing.