

Ecological site R108XD914MO

Wet Terrace Prairie

Last updated: 10/17/2024

Accessed: 05/20/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 108X—Illinois and Iowa Deep Loess and Drift

The Illinois and Iowa Deep Loess and Drift, Western Part MLRA covers parts of both Iowa and Missouri and is known locally as part of the Southern Iowa Drift Plain. A silty loess deposit of varying thickness (5 to 20 feet) covers a series of glacial advances known collectively as pre-Illinoian till. This till, deposited more than half a million years ago, was subjected to multiple instances of extreme erosion as well as periods of subdued erosion and intense weathering. The loess is thickest in the western part of the MLRA and generally thins eastward. In some areas, the loess has been removed and the older weathered till, called a “paleosol,” entirely exposed. These highly weathered soils, or paleosols, have a high content of clay, which slows the downward movement of water through the profile and causes water to move laterally instead of vertically. Wet areas, or “side-hill seeps,” commonly form where these paleosols become exposed along hillsides (Prior, 1991).

The dominant soil orders in this MLRA are Mollisols and Alfisols and, to a lesser extent, Entisols and Inceptisols. Most of the soils are Udolls or Udalfs. Aquolls are on the flatter interfluvies. The soils in the area dominantly have a mesic soil temperature regime, an aquic or udic soil moisture regime, and mixed mineralogy. They generally are very deep, well drained to poorly drained, and silty, loamy, or clayey. These soils on uplands include somewhat poorly drained, nearly level Argiudolls (Macksburg series); moderately well

drained, gently sloping to strongly sloping Argiudolls (Sharpsburg series); poorly drained, nearly level Argiaquolls (Winterset series); and well drained strongly, sloping to steep Hapludalfs (Gara, Lindley, Ladoga, and Armstrong series) (USDA-NRCS, 2006). The western part of the Illinois and Iowa Deep Loess and Drift is a segment of three other MLRAs within the Central Feed Grains and Livestock Region. The other areas are: the West-Central part (108C), the East-Central part (108B) and the Eastern part (108A).

Classification relationships

Major Land Resource Area (MLRA): Illinois and Iowa Deep Loess and Drift, Western Part (108D)

USFS Subregions: Central Dissected Till Plains Section (251C); Loess Hills (251Cb) and Central Dissected Till and Loess Plain (251Cc) Subsections (Cleland et al, 2007)

Relationship to Other Established Classifications:

NatureServe Classification: Ecological System: North-Central Interior Floodplain (9338);

Ecological Association: Central Wet-Mesic Tallgrass Prairie (NatureServe, 2013)

Landfire Biophysical Setting: Central Tallgrass Prairie (4314210) (Landfire, 2009)

Ecological site concept

Wet Terrace Prairies are within the red areas on the map (Figure 1). These sites formed in alluvium parent material and can be found on stream terraces on flood plains. Typically these sites are located along streams below loamy terrace savanna ecological sites. Soils are typically Mollisols, characterized by deep, dark colored surfaces high in organic matter due to the dominant prairie vegetation and have no rooting restrictions. Plant communities consist of mostly grasses and few forbs and shrubs

Associated sites

R108XD941IA	Loamy Terrace Savanna Loamy Terrace Savanna. Fine-silty soils including Watkins and Wiota series.
R108XD841IA	Loamy Foothlope Savanna Loamy Foothlope Savanna. Fine-loamy and fine-silty textured soils including Arbor, Ely, Judson and Olmitz series.
R108XD904IA	Wet Floodplain Prairie Wet Floodplain. Fine, fine-silty and fine-loamy soils including Ackmore, Amana, Carlow, Chequest, Colo, Lawson, Mt. Sterling, Sawmill, Spillville, Vesser, Wabash, Zook and Coland series.

Similar sites

R108XD941IA	Loamy Terrace Savanna Loamy Terrace Savanna. Fine-silty soils including Watkins and Wiota series.
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

Wet Terrace Prairies are of small extent, and can be found on stream terraces on flood plains along streams throughout MLRA 108D. Slopes are generally less than 5 percent. These sites typically occur on treads and risers of low stream terraces.




Figure 2. Block diagrams representing typical soil-landform sequences in Loess Ridges, Glacial Till Side/Footslopes. Red asterisk identifies soil component correlated to Wet Terrace Prairies.

Table 2. Representative physiographic features

Landforms	(1) Stream terrace
Runoff class	Negligible to low
Flooding frequency	None to occasional
Ponding frequency	None
Elevation	152–427 m
Slope	0–5%

Water table depth	0–122 cm
Aspect	Aspect is not a significant factor

Climatic features

The soil temperature regime of MLRA 108D is classified as “mesic” where the mean annual soil temperature is between 46 and 59°F (Soil Survey Staff, 2014). The average freeze-free period of this ecological site is about 175 days, while the frost-free period is about 150 days (Table 2). Average annual precipitation is 32 inches, which includes rainfall plus the water equivalent from snowfall (Table 3). The average annual low and high temperatures are 39 and 60°F, respectively.

Table 3. Representative climatic features

Frost-free period (characteristic range)	134-143 days
Freeze-free period (characteristic range)	157-172 days
Precipitation total (characteristic range)	889-940 mm
Frost-free period (actual range)	129-150 days
Freeze-free period (actual range)	150-180 days
Precipitation total (actual range)	889-940 mm
Frost-free period (average)	139 days
Freeze-free period (average)	164 days
Precipitation total (average)	914 mm

Climate stations used

- (1) GREENFIELD [USC00133438], Greenfield, IA
- (2) INDIANOLA 2W [USC00134063], Indianola, IA
- (3) WINTERSET 1N [USC00139132], Winterset, IA
- (4) BEDFORD [USC00130576], Bedford, IA
- (5) GUTHRIE CTR [USC00133509], Guthrie Center, IA
- (6) CRESTON 2 SW [USC00131962], Creston, IA
- (7) DES MOINES INTL AP [USW00014933], Des Moines, IA

Influencing water features

This site is poorly or somewhat poorly drained. Permeability is moderately slow to moderately rapid. The soil at this site is in hydrologic group C, D, or C/D (Hydrologic Soil Group, 2016). Land capability class is 1, 2e, 2w or 3w (Land Capability Classification, 2016). Depth of endosaturation is between 0 and 4 feet. This ecological site is influenced by a seasonal high water table from high groundwater levels, as well as slow hydraulic

conductivity, which impedes throughflow from precipitation and flood events. The water table is typically near the surface in late fall through spring, receding in the summer. This ecological site is on stream terraces and floodplain steps of perennial streams. They are not adjacent to the current stream channel. Areas on floodplain steps are subject to flooding, typically of short duration and low intensity. Constructed levees, often accompanied by stream channelization, have altered the flooding dynamics in many places.

Sites on floodplain steps are in the RIVERINE wetlands class of the Hydrogeomorphic (HGM) classification system (Brinson, 1993), whereas stream terrace areas are best considered as MINERAL SOIL FLAT wetlands. These areas are Emergent Palustrine wetlands (Cowardin et al., 1979).

Soil features

These soils have no major rooting restriction. The soils were formed under prairie vegetation, and have dark, organic-rich surface horizons. Parent material is alluvium. The soils have silt loam or silty clay loam surface horizons (Table 5). Subsoils are silt loam, silty clay loam or silty clay. Soil series associated with this site include Bremer, Humeston, Koszta, Nevin and Tuskeego.




Figure 9. Distribution of Wet Terrace Prairies within MLRA 108D

Table 4. Representative soil features

Parent material	(1) Alluvium
Surface texture	(1) Silt loam (2) Silty clay loam
Drainage class	Poorly drained to somewhat poorly drained

Permeability class	Very slow to moderately rapid
Soil depth	203 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (Depth not specified)	18.54–21.34 cm
Calcium carbonate equivalent (Depth not specified)	0%
Soil reaction (1:1 water) (Depth not specified)	5.5–6.5
Subsurface fragment volume <=3" (Depth not specified)	0%
Subsurface fragment volume >3" (Depth not specified)	0%

Ecological dynamics


Reference plant community is categorized as a wet mesic prairie and includes grasses and sedges, forbs, and shrubs. Species composition typically includes *Andropogon gerardii*, *Panicum virgatum*, *Spartina pectinate*, *Calamagrostis Canadensis*, *Helianthus grosseserratus*, *Juncus interior*, *Carex* spp., *Pycnanthemum virginianum*, *Oenothera pilosella*, and *Penstemon digitalis*.

Fire, grazing, drought, are all disturbances influencing the dynamics at this site. These sites likely burned 3 to 5 times every 10 years. Grazing by whitetail deer, prairie elk, was common, and bison as well may have been also present grazing to a lesser extent. Disturbances from these animals removed thatch, litter, and reduced the proliferation of small trees and shrubs (Mutel, 2008).

As this region was settled, these prairies were typically altered to better suit agricultural needs. Corn and soybean production is common in these areas today. Very few small remnants of this prairie exist now.

State and transition model

R108DY914MO Wet Terrace Prairie



Code	Process
T1A	Fire Suppression > 20 years; woody invasion
T1B	Tillage; vegetative seeding; grassland management
T1C, T3A, T5A	Tillage; conservation cropping system
T2A	Woody removal; tillage; vegetative seeding; grassland management
T2B	Woody removal; tillage; conservation cropping system
T4A	Vegetative seeding; grassland management
T3B, T4B	Vegetative seeding; prescribed fire; grassland management
1.1A	Fire-free interval 10+ years
1.2A	Fire interval 1-3 years
2.1A	Woody invasion; fire-free interval 40+ years
R2A	Woody removal; prescribed fire 1-3 years
R5A	Vegetative seeding; prescribed fire 1-3 years

Reference

As wet mesic prairie, this state has a reference plant community which is categorized as prairie and includes grasses, forbs and sedges. Periods of 10 years with no fire can cause this state to shift into a shrub invaded prairie (Figure 3). Conversely, fire intervals of 1 to 3 years will shift this phase back towards the reference community. Fire suppression greater than 20 years will cause this state to shift to a Woody Invaded Woodland State.

Restoration to the reference state is possible through removal of woody species and prescribed fires every 1-3 years. Conversion to cropland, or pasture are also typical transitions from reference state, the transition to cropland involves tillage and a conventional cropping system, and the transition to pasture is similar, requiring tillage, vegetative seeding, and grassland management.

Dominant plant species

- big bluestem (*Andropogon gerardii*), grass
- switchgrass (*Panicum virgatum*), grass
- prairie cordgrass (*Spartina pectinata*), grass

Community 1.1

Wet Mesic Tallgrass Prairie

A wet grassland ecosystem with numerous native species.

Dominant plant species

- big bluestem (*Andropogon gerardii*), grass
- switchgrass (*Panicum virgatum*), grass
- prairie cordgrass (*Spartina pectinata*), grass

Community 1.2

Shrub Invaded Prairie

Increase in woody species.

Dominant plant species

- big bluestem (*Andropogon gerardii*), grass
- switchgrass (*Panicum virgatum*), grass
- sawtooth sunflower (*Helianthus grosseserratus*), other herbaceous

Pathway P1.1A

Community 1.1 to 1.2

Fire free interval 10 plus years.

Pathway P1.2A

Community 1.2 to 1.1

Fire interval 1-3 Years

State 2

Woody Invaded Woodland

This woody invaded woodland forms as a result of a fire suppression interval of greater than 20 years on the reference state. The woody species have invaded enough to cause significant canopy closure. Restoration to the reference state requires removal of the woody species and a prescribed fire interval of 1 to 3 years. If the fire free period is 40 or more years, a closed woodland forms. Two transitions to other states are also possible. The transition to a cool season pasture state is accomplished through woody species removal, tillage, vegetative seeding and grassland management processes. The cropland state is the other possibility, requiring woody removal, tillage, and a conventional cropping system (Woodland Health, 2004).

Dominant plant species

- bur oak (*Quercus macrocarpa*), tree
- swamp white oak (*Quercus bicolor*), tree
- green bulrush (*Scirpus atrovirens*), grass

Community 2.1

Open Oak woodland

An increase in return fire interval results in a community with more woody species.

Dominant plant species

- bur oak (*Quercus macrocarpa*), tree
- swamp white oak (*Quercus bicolor*), tree
- green bulrush (*Scirpus atrovirens*), grass

Community 2.2

Closed Oak Woodland

longer fire intervals - increase in woody species

Dominant plant species

- bur oak (*Quercus macrocarpa*), tree
- swamp white oak (*Quercus bicolor*), tree
- slippery elm (*Ulmus rubra*), shrub
- Canadian blacksnakeroot (*Sanicula canadensis*), other herbaceous

Pathway P2.1A

Community 2.1 to 2.2

Woody invasion; fire free interval 40 plus years.

State 3

Non-Native Pasture

This state is formed from a native reference state, woody invaded woodland, or cropland which has been transformed into a cool season pasture due to several processes. In order to transform a native reference state, it requires tillage, vegetative seeding, and grassland management. From a woody invaded woodland, in addition to those processes involved in the reference state transition, it also requires woody removal. The Cropland transition to this state can be accomplished by only vegetative seeding and grassland management. Conversely, a transition to a cropland state from this state requires tillage and a conventional cropping system. This state can also transition to a native warm season grassland state by vegetative seeding, prescribed fire and grassland management processes.

Dominant plant species

- smooth brome (*Bromus inermis*), grass
- red clover (*Trifolium pratense*), other herbaceous

Community 3.1

Non-native pasture

seeded non-native grasses and forbs

Dominant plant species

- smooth brome (*Bromus inermis*), grass
- red clover (*Trifolium pratense*), other herbaceous

State 4

Cropland

In this state, tillage, seeding and herbicide has destroyed all of the original prairie. All other states can transition to this state through a combination of woody removal, if necessary, along with tillage, and a conventional tillage cropping system. Corn and soybeans are the principal crops. Variation in management within this state creates a wide range of soil properties and can be detrimental to the environment. Transitions to either a cool season pasture or a native warm season grassland are possible. The transition to cool season pasture state requires vegetative seeding and grassland management. The native warm season grassland state can be accomplished by vegetative seeding, prescribed fire and

grassland management.

Community 4.1

Conventional Tillage Field

Corn - soybean rotation is the most common crop.

State 5

Native Warm Season Grassland

The native warm season grassland state is a result of a transition from either a cool season pasture or cropland. Both require vegetative seeding, prescribed fire, and grassland management. It is possible to restore this state to the reference state by vegetative seeding, and a prescribed fire interval of 1 to 3 years.

Dominant plant species

- big bluestem (*Andropogon gerardii*), grass
- switchgrass (*Panicum virgatum*), grass
- Indiangrass (*Sorghastrum nutans*), grass

Community 5.1

Reconstructed prairie

Native warm season grass reconstructed prairie with various native forbs

Dominant plant species

- big bluestem (*Andropogon gerardii*), grass
- Indiangrass (*Sorghastrum nutans*), grass
- switchgrass (*Panicum virgatum*), grass

Transition T1A

State 1 to 2

Fire suppression of 20 years or more; woody invasion.

Transition T1B

State 1 to 3

Tillage; vegetative seeding; grassland management

Transition T1C

State 1 to 4

Tillage; conservation cropping system.

Restoration pathway R2A

State 2 to 1

Woody removal; prescribed fire 1-3 years.

Transition T2A

State 2 to 3

Woody removal; tillage; vegetative seeding; grassland management

Transition T2B

State 2 to 4

Woody removal; tillage; conservation cropping system.

Transition T3A

State 3 to 4

Tillage; conservation cropping system.

Transition T3B

State 3 to 5

Vegetative seeding; prescribed fire; grassland management.

Restoration pathway T4A

State 4 to 3

Vegetative seeding; grassland management

Transition T4B

State 4 to 5

Vegetative seeding; prescribed fire; grassland management.

Restoration pathway R5A

State 5 to 1

Vegetative seeding; prescribed fire 1-3 years.

Transition T5A

State 5 to 4

Tillage; conservation cropping system

Additional community tables

Inventory data references

No field plots were available for this site. A review of the scientific literature and professional experience were used to approximate the plant communities for this provisional ecological site. Information for the state-and-transition model was obtained from the same sources. All community phases are considered provisional based on these plots and the sources identified in ecological site description.

Other references

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C. Carpenter, and W.H. McNab. 2007. Ecological Subregions: Sections and Subsections of the Conterminous United States. USDA Forest Service, General Technical Report WO-76. Washington, DC.

Landfire. 2009. Biophysical Setting 4314210 Central Tallgrass Prairie. In: Landfire National Vegetation Dynamics Models. USDA For. Serv. and U.S. Department of Interior. Washington, DC.

Mutel, Cornelia F. 2008. The Emerald Horizon: The History of Nature in Iowa. University of Iowa Press.

NatureServe. 2013. Associations and alliances of Iowa. St. Paul, Minnesota.

Prior, Jean Cutler. 1991. Landforms of Iowa. University of Iowa Press. Iowa City, Iowa.

Soil Survey Staff. 2014. Keys to Soil Taxonomy, 12th ed. USDA-Natural Resources Conservation Service, Washington, DC.

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. USDA Handbook 296.

United States Department of Agriculture, Natural Resources Conservation Service. Hydrologic Soil Group. National soil survey handbook, title 430-VI.
http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ref/?cid=nrcs142p2_054223#39
accessed August 29, 2016.

United States Department of Agriculture, Natural Resources Conservation Service. Land Capability Classification. National soil survey handbook, title 430-VI.
http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ref/?cid=nrcs142p2_054226#02

accessed August 29, 2016.

Woodland Health: Stewardship Options for Iowa Woodland Owners. 2004. Iowa Natural Heritage Foundation. Des Moines, Iowa.

Contributors

John Hammerly (john.hammerly@usda.gov)

Dan Pulido (dan.pulido@usda.gov)

Approval

Suzanne Mayne-Kinney, 10/17/2024

Acknowledgments

This ESD was originally approved prior to April 2021.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	John Hammerly, soil scientist Dan Pulido, SSOL
Contact for lead author	
Date	05/20/2025
Approved by	Suzanne Mayne-Kinney
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater**

than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

- 14. Average percent litter cover (%) and depth (in):**
-

- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

- 17. Perennial plant reproductive capability:**
-